

# Widely Distributed Mutations in the COL2A1 Gene Produce Achondrogenesis Type II/Hypochondrogenesis

Jarmo K  rkk  ,<sup>1,2</sup> Daniel H. Cohn,<sup>3,5</sup> Leena Ala-Kokko,<sup>1,2</sup> Deborah Krakow,<sup>3,4,6</sup> and Darwin J. Prockop<sup>1\*</sup>

<sup>1</sup>Center for Gene Therapy, MCP Hahnemann University, Philadelphia, Pennsylvania

<sup>2</sup>Collagen Research Unit, Biocenter Oulu, and Department of Medical Biochemistry, University of Oulu, Oulu, Finland

<sup>3</sup>Ahmanson Department of Pediatrics, Steven Spielberg Pediatric Research Center, Los Angeles, California

<sup>4</sup>Department of Obstetrics and Gynecology, Burns and Allen Cedars-Sinai Research Institute, Los Angeles, California

<sup>5</sup>Departments of Pediatrics, UCLA School of Medicine, Los Angeles, California

<sup>6</sup>Department of Obstetrics and Gynecology, UCLA School of Medicine, Los Angeles, California

The COL2A1 gene was assayed for mutations in genomic DNA from 12 patients with achondrogenesis type II/hypochondrogenesis. The exons and flanking sequences of the 54 exons in the COL2A1 gene were amplified by a series of specific primers using PCR. The PCR products were scanned for mutations by conformation sensitive gel electrophoresis, and PCR products that generated heteroduplex bands were then sequenced. Mutations in the COL2A1 gene were found in all 12 patients. Ten of the mutations were single base substitutions that converted a codon for an obligate glycine to a codon for an amino acid with a bulkier side chain. One of the mutations was a change in a consensus RNA splice site. Another was an 18-base pair deletion of coding sequences. The results confirmed previous indications that conformation sensitive gel electrophoresis is highly sensitive for detection of mutations in large and complex genes. They also demonstrate that most, if not all, patients with achondrogenesis type II/hypochondrogenesis have mutations in the COL2A1 gene. *Am. J. Med. Genet.* 92:95–100, 2000.   2000 Wiley-Liss, Inc.

**KEY WORDS:** achondrogenesis; hypochondrogenesis; osteochondrodysplasia; COL2A1; collagen

## INTRODUCTION

Type II collagen is the most abundant protein of the cartilage matrix and is essential for the normal embryonic development of the skeleton, for linear growth, and for the ability of cartilage to resist compressive forces [Piez, 1984; Prockop and Kivirikko, 1995]. Mutations in the gene for type II collagen, COL2A1, disrupt these processes and cause a variety of osteochondrodysplasias [Ritvaniemi et al., 1995; Kuivaniemi et al., 1997]. COL2A1 mutations were first identified in patients with spondyloepiphyseal dysplasia and hypochondrogenesis [Lee et al., 1989; Vissing et al., 1989]. Since then, over 40 mutations in the COL2A1 gene have been reported in patients with achondrogenesis type II/hypochondrogenesis (ACG2/HCG), spondyloepimetaphyseal dysplasia Strudwick type, Kniest dysplasia, the Stickler and Wagner syndromes, and mild osteochondrodysplasia phenotypes with early-onset osteoarthritis as a major manifestation [Kuivaniemi et al., 1997; Spranger et al., 1994]. Most of the mutations were single base changes that each caused a single amino acid substitution for one of the obligate glycine residues that appear in the repeating Gly-X-Y triplet characteristic of collagen  $\alpha$ -chains. However, RNA splicing mutations, partial gene deletions, insertions, and duplications were also observed. In addition, five families, three of whom perhaps were related through an early ancestor, had a single base mutation that changed the codon for arginine at residue 519 to a cysteine codon (R519C) and produced primary generalized osteoarthritis with or without evidence of a mild osteochondrodysplasia phenotype [Ala-Kokko et al., 1990; Pun et al., 1994; Williams et al., 1995; Bleasel et al., 1995; Holderbaum et al., 1996]. In three unrelated

Grant sponsor: National Institutes of Health; Grant numbers: HD22657, HD01205; Grant sponsor: Arthritis Foundation; Grant sponsor: Finnish Cultural Foundation.

\*Correspondence to: Dr. Darwin J. Prockop at the Center for Gene Therapy, MCP Hahnemann University, 10118NCB, Mail Stop 421, 245 North 15th Street, Philadelphia, PA 19102. E-mail prockop@mcphu.edu

Received 5 March 1999; Accepted 17 December 1999

TABLE I. Clinical Data on Patients

ISDR <sup>a</sup> number	Gestation age (weeks)	Gender	Radiograph diagnosis	Cartilage morphology	Hydrops	Midface hypoplasia	Cleft palate	Other abnormalities
85-155	35	F	ACG2	Hypervascular, hypercellular	+	+	+	
87-051	23	M	ACG2	Sparse collagen fibrils, hypervascular, large foamy lacunae	+	+	—	Polyhydramnios
89-123	30	F	ACG2	Sparse matrix, hypervascular, chondrocyte RER inclusions	+	+	+	Polyhydramnios
89-131	22	F	ACG2	Sparse matrix, hypervascular, chondrocyte RER inclusions	+	+	Unknown	
89-200	36	F	ACG2	Hypervascular, chondrocyte RER inclusions	+	+	+	Polyhydramnios
91-215	24	M	ACG2	Sparse matrix, hypervascular, chondrocyte RER inclusions	+	+	Unknown	Polyhydramnios
92-197	21	M	ACG2/HCG	Sparse matrix, hypervascular	+	Unknown	Unknown	
92-240	15	F	ACG2	Hypervascular, hypercellular cartilage	+	+	+	Cystic hygroma
93-078	23	F	ACG2	Unknown	—	+	—	Polyhydramnios
93-260	28	F	ACG2	Foamy lacunae	+	+	+	Polyhydramnios, dysplastic left kidney, bilateral equinovarus
94-160	17	F	ACG2	Hypervascular, hypercellular	+	+	Unknown	Bilateral equinovarus
95-085	32	F	ACG2	Sparse matrix, hypercellular, foamy lacunae, chondrocyte RER inclusions, disorganized growth plate	+	+	Unknown	Polyhydramnios, cystic hygroma, hydrocephalus

<sup>a</sup>International Skeletal Dysplasia Registry.

patients, a R75C substitution produced a similar phenotype [Bleasel et al., 1995; Williams et al. 1993]. In contrast, a R789C substitution, identified in two unrelated patients, resulted in spondyloepiphyseal dysplasia [Chan et al., 1990, 1993].

Although a large number of mutations in the COL2A1 gene have been reported in patients with varying phenotypes, it has been difficult to define the true incidence of such mutations among patients with a specific phenotype [Ritvaniemi et al., 1995]. This is primarily because patient cartilage is usually not available for analysis at the protein or mRNA level, and the COL2A1 gene has a complex structure composed of 54 exons. Thus complete examination of the COL2A1 gene for mutations is technically difficult. The highest number of mutations reported in the COL2A1 gene was in patients with the perinatally lethal phenotypes of ACG2 and HCG, with 13 reported mutations [Ritvaniemi et al., 1995; Kuivaniemi et al., 1997; Mundlos et al., 1996]. To determine the sensitivity of the relatively simple technique [Ganguly et al., 1993] of conformation sensitive gel electrophoresis (CSGE) for detecting COL2A1 mutations, and to examine the spectrum of mutations that can produce ACG2/HCG, we carried out mutation analysis for 12 patients. Mutations were identified in all 12 patients, demonstrating that CSGE is highly sensitive for detecting mutations in the COL2A1 gene, that ACG2/HCG appears to be produced exclusively by COL2A1 mutations, and that the mutations are distributed across the gene more widely than previously assumed.

## MATERIALS AND METHODS

### Clinical Ascertainment

Twelve fetuses with ACG2 or HCG were ascertained through the International Skeletal Dysplasia Registry

(ISDR). The gestational ages of the fetuses ranged from 15 to 35 weeks. Clinically the fetuses presented with findings typical for the ACG2/HCG spectrum: hydrops, midface hypoplasia, cleft palate, severe shortness of the rhizo- and meso-melic portions of the limbs, and




Fig. 1. AP view of fetus R93-260.

equinovarus (summarized in Table I) [Chen et al., 1981]. The radiographs were also consistent with the diagnosis, demonstrating severely underossified vertebral bodies, small iliac bones, shortened tubular bones with metaphyseal cupping, lateral bowing of the femurs, and unossified pubic and ischial bones (Fig. 1).

Cartilage morphology was examined in 11 of the cases (see Table I) by published methods [Brodie et al., 1998]. In all cases, the characteristic findings of hypervascular/hypercellular cartilage, foamy lacunae, a sparse matrix, and rough endoplasmic reticulum inclusions in chondrocytes (Fig. 2) were also consistent with the diagnosis of ACG2/HCG [Horton et al., 1987].

### Mutation Analysis

Genomic DNA was extracted from blood samples or from cultured skin fibroblasts. The exons and the flanking sequences of the 54 exons of the COL2A1 gene were amplified by a series of specific primers (<http://www.mcphu.edu/medschool/centers/matrix/images/col2a1pr.htm>) based on the published gene sequence [Ritvaniemi et al., 1995; Williams et al., 1993, 1995; Ala-Kokko and Prockop, 1990]. Genomic DNA was amplified in a 40  $\mu$ l volume by thermal cycling at 95°C for 10 min for one cycle, followed by 95°C for 40 sec, 60°C for 40 sec, and 72°C for 50 sec, for 35 cycles. This was




Fig. 2. Transmission electron micrograph of cartilage from fetus R95-085. Final magnification: 20,000 $\times$ .



followed by heteroduplex formation steps: 95  C for 5 min and 68  C for 30 min. CSGE analysis was performed as described elsewhere [Ganguly et al., 1993, K  rkk   et al., 1998a,b]. The gels were examined with a hand-held UV illuminator (short wave) to identify regions containing homoduplex and heteroduplex fragments. Appropriate regions were excised from the gels, transferred to filter paper, and analyzed further in a UV-image analyzer with a CCD camera (DOC-IT<sup>TM</sup> Gel Documentation System; UVP). The image from the monitor was recorded by a thermal printer. After treatment with exonuclease I and shrimp alkaline phosphatase, the sequences of PCR products containing heteroduplexes were determined by using the ABI PRISM Dye Terminator Cycle Sequencing Ready Kit with AmpliTaq DNA polymerase, FS (Perkin-Elmer, Foster City, CA) and an automated instrument (ABI PRISM 377 Sequencer; Perkin-Elmer). PCR products that contained deletions in one allele were purified from agarose (QIA II Gel Extraction Kit; Qiagen, Chatsworth, CA) and cloned into a plasmid (pT7 Blue T-Vector Kit; Novagen, Madison, WI) prior to sequence analysis.

## RESULTS

All 54 exons and the flanking sequences of the COL2A1 gene were amplified by PCR from all 12 patients with ACG2/HCG. The samples were then analyzed for mutations in the COL2A1 gene. Mutations were first identified by unique heteroduplexes detected by CSGE (Fig. 3), and sequence analysis of the PCR products was used to define the specific mutations. The analysis identified novel disease-causing mutations in all patients (Table II). Ten of the mutations were substitutions for obligate glycines in the triple helical domain of the type II collagen molecule, one was a deletion of 18 bp of coding sequence, and one was predicted to cause exon skipping. In addition to the disease-causing mutations, over 30 neutral polymorphisms were detected (Table III).




Fig. 3. The CSGE patterns caused by the mutations. Numbers indicate the patient numbers. C, Control. Lanes 1–12 correspond to patients ISDG 89–131 to 94–160 in Table II.

TABLE II. Mutations in COL2A1 Causing Achondrogenesis II/Hypochondrogenesis

ISDG number	Patient	Mutation in COL2A1	
		Exon	Mutation
89-131	ACG2	19	G253V
89-200	ACG2	19	G253D
85-155	HCG	32	IVS34 + 1G > A
92-240	ACG2	33	G571D
87-051	ACG2	33	G580R
91-215	ACG2	34	G595R
92-197	ACG2/HCG	39	G694E
95-085	ACG2	40	G748D
89-123	ACG2	41	G781S
93-078	HCG	44	G817-V822del6aa
93-260	ACG2	44	G865V
94-160	ACG2	46	G919R

## DISCUSSION

On phenotypic, radiographic, and morphologic grounds [Chen et al., 1981; Horton et al., 1987], as well as in the light of mutation analyses of the COL2A1 gene, it was previously assumed that achondrogenesis type II/hypochondrogenesis was caused exclusively by COL2A1 mutations. However, because of limitations in the sensitivity of mutation detection methods employed, no systematic analysis of patients was possible. Furthermore, for collagen genes, including COL2A1, the large number of highly homologous exons has presented a formidable technical barrier for the complete detection of mutations [Prockop and Kivirikko, 1995].

TABLE III. Polymorphisms Detected in the COL2A1 Gene

Fragment for exon number	Change
1	TCG-ACG (SerN9-Thr)
1	IVS1+18g/c
5a	GAA-GAT (GluN142)
5b	GGC-GGA (GlyN168)
6	IVS5-27t/a
7	IVS6-7g/a
7	IVS7+15g/t
9	IVS9+15g/a
9	IVS9+42delg
10	IVS9-42g/a
18	IVS18+7g/a
20	IVS19-11c/t
30	IVS29-49g/t
30	IVS30+7a/g
30	IVS30+12g/c
32	IVS31-53 <sup>^</sup> -52insg
32	GGC-GGT (Gly565)
32	IVS32+73c/t
33	IVS32-32t/c
33	IVS32-22g/a
33	IVS33+68g/a
34	IVS33-56g/t
34	AAC-AAT (Asn600)
34	IVS34+84c/g
36	IVS35-74a/c
40	IVS39-45delg
50	GTT-ATT (Val1131Ile)
50	IVS50+48-+49delct
51	IVS50-14c/g
52	3' UTR+4c/t




Fig. 4. Schematic of the type II collagen molecule indicating sites of mutation causing achondrogenesis type II/ hypochondrogenesis. Arrows above molecule: the sites of mutations reported here. Arrows below molecule: the sites of previously reported mutations. <sup>1</sup>The mutation detected was IVS34+1G→A; the figure presents the predicted effect on the amino acid level. <sup>2</sup>The exact amino acid number not known [Mundlos et al., 1996].

The use of CSGE [Ganguly et al., 1993, 1997; Gatalica et al., 1997; Körkkö et al. 1998a,b; Philippe et al., 1997] has greatly facilitated the analysis of such genes because it has provided a relatively simple means of scanning PCR products for single base mismatches or other allelic differences. The procedure has been shown to be sensitive enough to detect essentially all single base changes in several hundred alleles of the COL1A1, COL1A2, and COL2A1 genes [Ganguly et al., 1993; Körkkö et al., 1998b]. The results presented here further demonstrate the sensitivity of this analytic method and suggest that the phenotypes studied are caused primarily, if not exclusively, by defects in type II collagen.

All 13 previously reported mutations in the ACG2/HCG patients caused conversion of an obligate glycine to a bulkier amino acid [Kuivaniemi et al., 1997; Mundlos et al., 1996]. The data presented here demonstrate that in-frame deletions and RNA splicing mutations can also produce phenotypes at the lethal end of the type II collagenopathy spectrum. Based on a similar hypothesis that emerged from analysis of the type I collagen genes in patients with osteogenesis imperfecta [Byers, 1993], it was suggested that type II collagen gene mutations in patients with more severe phenotypes might cluster at the carboxyl-terminal end of the triple helical domain [Murray et al., 1989]. This hypothesis was consistent with mutations previously identified in ACG2/HCG since, with the exception of a G310D substitution characterized in a case of ACG2 [Bonaventure et al., 1995], all mutations were located in the region of the gene corresponding to the carboxyl-terminal half of the triple helical domain. Mutations in the cohort studied here demonstrate that lethal COL2A1 mutations are more widely distributed in the molecule, a result not consistent with the gradient hypothesis (Fig. 4).

Studies with synthetic peptides [Chan et al., 1997], as well as genotype/phenotype comparisons in patients [see Westerhausen et al., 1990; Wang et al., 1993; Marini et al., 1993], demonstrate that the sequence context and the specific amino acid substitutions for glycine influence the functional consequences. However, it has been difficult to establish firm relationship between the mutation and the severity of the phenotype because there are several examples of the same substitution of an amino acid for glycine at the same position that produce phenotypes that vary from mild or moderate to lethal (Kuivaniemi et al., 1997). The phenotypic variability seen with similar or identical mutations in collagen genes is consistent with the phenotype variability recently seen in monozygotic twins with identical mutations in a variety of genes [see Hall, 1996; Wyszyski et al., 1996; Yamagishi et al., 1998].

#### ACKNOWLEDGMENTS

We thank the families and their health care providers who referred the cases to the International Skeletal Dysplasia Registry. We thank the members of the Registry, D.L. Rimo, R.S. Lachman, M. Priore, and R. Bonaquisti for their assistance with diagnostic and organizational efforts. We thank L. Nolasco, B. Mekikian, H.E. Gruber, and W.R. Wilcox for morphologic studies. We thank Mr. Robert Hnatuk and Mr. Richard Kosich for expert technical assistance. This work was supported in part by grants HD22657, and HD01205 from the National Institutes of Health, a grant from the Arthritis Foundation (DJP) and a grant from Finnish Cultural Foundation (JK).

#### REFERENCES

- Ala-Kokko L, Baldwin CT, Moskowitz RW, Prockop DJ. 1990. Single base mutation in the type II procollagen gene (COL2A1) as a cause of pri-

- mary osteoarthritis associated with a mild chondrodysplasia. *Proc Natl Acad Sci USA* 87:6565–6568.
- Ala-Kokko L, Prockop DJ. 1990. Completion of the intron-exon structure of the gene for human type II procollagen (COL2A1): variations in the nucleotide sequence of the alleles from three chromosomes. *Genomics* 8:454–460.
- Bleasel JF, Holderbaum D, Haqqi TM, Moskowitz RW. 1995. Clinical correlations of osteoarthritis associated with single base mutations in the type II procollagen gene. *J Rheumatol* 43:34–36.
- Bonaventure J, Cohen-Solal L, Ritvaniemi P, Van Maldergem L, Kadhom N, Delezoide AL, Maroteaux P, Prockop DJ, Ala-Kokko L. 1995. Substitution of aspartic acid for glycine at position 310 in type II collagen produces achondrogenesis II, and substitution of serine at position 805 produces hypochondrogenesis: analysis of genotype-phenotype relationships. *Biochem J* 307:823–30.
- Brodie SG, Lachman RS, Crandall BF, Fox MA, Rimoin DL, Cohn DH, Wilcox WR. 1998. Radiographic and morphologic findings in a previously undescribed type of mesomelic dysplasia resembling atelosteogenesis type II. *Am J Med Genet* 80:247–51.
- Byers PH. 1993. Osteogenesis imperfecta. In: Royce PM, Steinmann B, editors. *Connective tissue and its heritable disorders*. New York: Wiley-Liss. p 317–351.
- Chan D, Rogers JF, Bateman JF, Cole WG, Ala-Kokko L, Prockop DJ. 1990. Recurrent substitutions of arginine 789 by cysteine in pro-alpha 1 (II) collagen chains produce spondyloepiphyseal dysplasia congenita. *J Rheumatol Suppl* 43:37–38.
- Chan D, Taylor TK, Cole WG. 1993. Characterization of an arginine 789 to cysteine substitution in alpha 1 (II) collagen chains of a patient with spondyloepiphyseal dysplasia. *J Biol Chem* 15:15238–15245.
- Chan VC, Ramshaw JAM, Kirkpatrick A, Beck K, Brodsky B. 1997. Positional preferences of ionizable residues in Gly-X-Y triplets of the collagen triple-helix. *J Biol Chem* 272:31441–31446.
- Chen H, Liu CT, Yang SS. 1981. Achondrogenesis: a review with special consideration of achondrogenesis type II (Langer-Saldino). *Am J Med Genet* 10:379–395.
- Ganguly A, Rock MJ, Prockop DJ. 1993. Conformation-sensitive gel electrophoresis for rapid detection of single-base differences in double-stranded PCR products and DNA fragments. evidence for solvent-induced bends in DNA heteroduplexes. *Proc Natl Acad Sci USA* 90:10325–10329.
- Ganguly A, Williams C. 1997. Detection of mutations in multiple-exon genes comparison of conformation sensitive gel electrophoresis and sequencing strategies with respect to cost and time for finding mutations. *Hum Mutat* 9:339–343.
- Gatalica B, Pulkkinen L, Li K, Kuokkanen K, Ryynänen M, McGrath JA, Uitto J. 1997. Cloning of the human type XVII collagen gene (COL17A1) detection of novel mutations in generalized atrophic benign epidermolysis bullosa. *J Hum Genet* 60:352–365.
- Hall JG. 1996. Twinning: mechanisms and genetic implications. *Curr Opin Genet Dev* 6:343–347.
- Holderbaum D, Bleasel JF, Johnsson H, Saxne T, Prockop DJ, Williams CJ, Moskowitz RW. 1996. Arg519-Cys-associated haplotype in Icelandic Family suggest early origin of this mutation. *Matrix Biol* 15:178.
- Horton WA, Machado MA, Chou JW, Campbell D. 1987. Achondrogenesis type II, abnormalities of the extracellular matrix. *Pediatr Res* 22:234–329.
- Körkkö J, Ala-Kokko L, De Paepe A, Nuytinck L, Earley J, Prockop DJ. 1998a. Analysis of the COL1A1 and COL1A2 genes by PCR amplification and scanning by conformation sensitive gel electrophoresis identifies only COL1A1 mutations in 15 patients with mild osteogenesis imperfecta (type I): identification of common sequences null allele mutations. *Am J Hum Genet* 62:98–110.
- Körkkö J, Annunen S, Pihlajamäa T, Prockop DJ, Ala-Kokko L. 1998b. Conformation sensitive gel electrophoresis (CSGE) for simple and accurate detection of mutations. Comparison with denaturing gradient gel electrophoresis (DGGE) and nucleotide sequencing. *Proc Natl Acad Sci USA* 95:1681–1685.
- Kuivaniemi H, Tromp G, Prockop DJ. 1997. Mutations in fibrillar collagens (types I, II, III and XI), fibril-associated collagen (type IX), and network-forming collagen (type X) cause a spectrum of diseases of bone, cartilage and blood vessels. *Hum Mutat* 9:300–315.
- Lee B, Vissing H, Ramirez F, Rogers D, Rimoin D. 1989. Identification of the molecular defect in a family with spondyloepiphyseal dysplasia. *Science* 26:978–980.
- Marini JC, Lewis MB, Wang Q, Chen KJ, Orrison BM. 1993. Serine for glycine substitutions in type I collagen in two cases of type IV osteogenesis imperfecta (OI). Additional evidence for a regional model of OI pathophysiology. *J Biol Chem* 268:2667–2673.
- Mundlos S, Chan D, Bateman JF. 1996. An I(II) Gly913 to Cys substitution prevents the matrix incorporation of type II collagen which is replaced with type I and III collagens in cartilage from a patient with hypochondrogenesis. *Am J Med Genet* 63:129–136.
- Murray LW, Bautista J, James PL, Rimoin DL. 1989. Type II collagen defects in the chondrodysplasias. I. Spondyloepiphyseal dysplasias. *Am J Hum Genet* 45:5–15.
- Philippe C, Porter DE, Emerton ME, Wells DE, Simpson AH, Monaco AB. 1997. Mutation screening of EXT1 and EXT2 genes in patients with hereditary multiple exostosis. *J Hum Genet* 61:520–528.
- Piez KA. 1984. Molecular and aggregate structures of collagens. In: Piez KA, Reddi AH, editors. *Extracellular matrix biochemistry*. New York: Elsevier Science Publishing Co. p 1–40.
- Prockop DJ, Kivirikko KI. 1995. Collagen: Molecular biology, diseases and potentials for therapy. *Annu Rev Biochem* 64:403–434.
- Pun YL, Moskowitz RW, Lie S, Sundstrom WR, Block SR, McEwen C, Williams HJ, Bleasel JF, Holderbaum D, Haqqi TM. 1994. Clinical correlations of osteoarthritis associated with a single-base mutation (arginine 519 to cysteine) in type II procollagen gene: a newly defined pathogenesis. *Arth Rheum* 37:264–9.
- Ritvaniemi P, Körkkö J, Bonaventure J, Vikkula M, Hyland J, Paasilta P, Kaitila I, Kääriäinen H, Sokolov BP, Hakala M, Mannismäki P, Meerson EM, Klemola T, Williams C, Peltonen L, Kivirikko KI, Prockop DJ, Ala-Kokko L. 1995. Analysis of the COL2A1 gene detects mutations in over 20 percent of patients with chondrodysplasias and up to 2 percent of patients with familial osteoarthritis. *Arth Rheum* 38:999–1004.
- Spranger J, Winterpacht A, Zabel B. 1994. The type II collagenopathies: a spectrum of chondrodysplasias. *Eur J Pediatr* 153:56–65.
- Vissing H, D'Alessio M, Lee B, Ramirez F, Godfrey M, Hollister DW. 1989. Glycine to serine substitution in the triple helical domain of proI(II) collagen results in a lethal perinatal form of short-limbed dwarfism. *J Biol Chem* 264:18265–18267.
- Wang Q, Orrison BM, Marini JC. 1993. Two additional cases of osteogenesis imperfecta with substitutions for glycine in the alpha 2(I) collagen chain: a regional model relating mutation location with phenotype. *J Biol Chem* 268:25162–25167.
- Westerhausen A, Kishi J, Prockop DJ. 1990. Mutations that substitute serine for glycine alpha 1-598 and glycine alpha 1-631 in type I procollagen: the effects on thermal unfolding of the triple helix are position-specific and demonstrate that the protein unfolds through a series of cooperative blocks. *J Biol Chem* 15:13995–4000.
- Williams CJ, Considine EL, Knowlton RG, Reginato A, Neumann G, Harrison D, Buxton P, Jimenez S, Prockop DJ. 1993. Spondyloepiphyseal dysplasia and precocious osteoarthritis in a family with an Arg75 to Cys mutation in the procollagen type II gene (COL2A1). *Hum Genet* 92:499–500.
- Williams CJ, Rock M, Considine E, McCarron S, Gow P, Ladda R, McLain D, Michels VM, Murphy W, Prockop DJ, Ganguly A. 1995. Three new point mutations in type II procollagen (COL2A1) and identification of a fourth family with the COL2A1 Arg519 to Cys base substitution using conformation sensitive gel electrophoresis. *Hum Mol Genet* 4:309–312.
- Wyszynski DF, Lewanda AF, Beaty TH. 1996. Phenotypic discordance in a family with monozygotic twins and non-syndromic cleft lip and palate. *Am J Med Genet* 66:468–470.
- Yamagishi H, Ishii C, Maeda J, Kojima Y, Matsuoka R, Kimura M, Takao A, Momma K, Matsuo N. 1998. Phenotypic discordance in monozygotic twins with 22q11.2 deletion. *Am J Med Genet* 78:319–321.